
1. Let *X* be a continuous random variable with pdf

$$f(x) = \begin{cases} x & 0 \le x \le 1, \\ 2 - x & 1 \le x \le 2, \\ 0 & \text{otherwise.} \end{cases}$$

(a) Sketch the pdf of *X*.



(b) Find the cdf of *X* and sketch it.


If 
$$0 \le x \le 1$$
, then  $F(x) = \int_{0}^{x} y \, dy = \frac{1}{2} y \Big|_{y=0}^{y=x} = \frac{1}{2} x^{2}$   
 $cdf: P(X \le x)^{n}$   $f(X \le x)^{n}$   
IF  $1 \le x \le 2$ , then  $F(x) = \frac{1}{2} + \int_{1}^{x} (2-y) \, dy = \frac{1}{2} + \left[ 2y - \frac{1}{2}y^{2} \right]_{y=1}^{y=x} = 2x - \frac{x^{2}}{2} - 1$   
 $P(X \le 1)^{n}$   $P(1 \le x \le x)^{3}$   
Thus:  
 $F(x) = \begin{cases} 0 & \text{if } x < 0 \\ \frac{1}{2}x^{2} & \text{if } 0 \le x \le 1 \\ 2x - \frac{x^{2}}{2} - 1 & \text{if } 1 < x \le 2 \\ 1 & \text{if } 2 < x \end{cases}$ 

(c) What is 
$$P(X < 1.5)$$
?  
Find the area under the pdf left of  $x = 1.5$ .  
 $P(X < 1.5) = \int_{-1}^{15} f(x) dx = \frac{7}{8}$ 

(d) Find a value  $\eta_{0.75}$  such that  $P(X \le \eta_{0.75}) = 0.75$ .

Find C such  $\frac{c^2}{2} = \frac{1}{4}$  so  $\eta_{0.75} = 2 - \frac{12}{2}$ that  $c \chi^{45^{\circ}}$   $c^2 = \frac{1}{2}$   $\approx 1.293$ to  $r_{0.75} = 1.293$  2. Suppose that a continuous random variable *X* has pdf f(x) = kx(4 - x) for  $0 \le x \le 4$ , and f(x) = 0 otherwise.

(a) Sketch the pdf of *X*. Then, without computing anything, sketch the cdf of *X*.



(b) What is the value of *k*?

Remember that the pdf must integrate to 1:  

$$\int_{0}^{4} kx (4-x) dx = \frac{32}{3} k = 1, \quad \text{so} \quad k = \frac{3}{32}.$$

(c) Find P(X > 3 or X < 1).

$$P(X > 3 \text{ or } X < 1) = \int_{0}^{1} f(x) dx + \int_{3}^{4} f(x) dx = 2 \int_{0}^{1} \frac{3}{32} x(4-x)$$
  
by symmetry of  $f(x)$  about  $x=2$   
$$= \frac{3}{16} \left[ 2x^{2} - \frac{x^{3}}{3} \right]_{0}^{1} = \frac{3}{16} \left( 2 - \frac{1}{3} \right) = \frac{5}{16}$$

- 3. Suppose that the cdf of a random variable *X* is  $F(x) = 1 e^{-5x}$  for x > 0, and F(x) = 0 otherwise.
  - (a) What is the pdf of *X*? Sketch both the pdf and the cdf.

Differentiate the cdf to find the pdf:  

$$f(x) = \frac{d}{dx}F(x) = \frac{d}{dx}(1 - e^{-5x}) = 5e^{-5x} \text{ for } x > 0.$$

$$f(x) = \frac{d}{dx}F(x) = \frac{d}{dx}(1 - e^{-5x}) = 5e^{-5x} \text{ for } x > 0.$$

- (b) What is  $P\left(\frac{1}{4} < X < \frac{1}{3}\right)$ ? Can you get this from *either* the cdf or the pdf?
  - $cdf: P\left(\frac{1}{4} < X < \frac{1}{3}\right) = F\left(\frac{1}{3}\right) F\left(\frac{1}{4}\right) = \left(1 e^{-\frac{5}{4}}\right) \left(1 e^{-\frac{5}{4}}\right) = e^{-\frac{5}{4}} e^{-\frac{5}{3}} \approx 0.098$


$$\mathsf{Pdf}: \qquad \mathsf{P}\left(\frac{1}{4} < X < \frac{1}{3}\right) = \int_{\frac{1}{4}}^{\frac{1}{3}} \mathsf{f}(x) dx = \mathsf{same}$$

4. Random variable *X* has pdf  $f(x) = \begin{cases} ax + bx^2 & 0 < x < 1, \\ 0 & \text{otherwise.} \end{cases}$ 

Furthermore, 
$$P\left(X < \frac{1}{2}\right) = \frac{3}{16}$$
. What is the median of X?  
We need:  $\int_{0}^{1} (ax + bx^{2}) dx = \frac{a}{2} + \frac{b}{3} = 1$   $\Rightarrow 3a + 2b = 6$   
 $\int_{0}^{\frac{1}{2}} (ax + bx^{2}) dx = \frac{a}{2} \cdot \frac{1}{4} + \frac{b}{3} \cdot \frac{1}{8} = \frac{a}{8} + \frac{b}{24} = \frac{3}{16} \Rightarrow 3a + b = \frac{9}{2}$   
Median:  $\int_{0}^{1} (x + \frac{3}{2}x^{2}) dx = \frac{1}{2}\eta^{2} + \frac{1}{2}\eta^{3} = \frac{1}{2} \Rightarrow \eta^{2} + \eta^{3} = 1$   
So we need  $\eta^{3} + \eta^{2} - 1 = 0$ , or  $\eta \approx 0.755$   
 $E_{xact}$ :  $\eta = \frac{1}{3} \left[ -1 + \sqrt[3]{\frac{25}{3} - \frac{3\sqrt{69}}{2}} + \sqrt[3]{\frac{1}{2}(25 + 3\sqrt{69})} \right]$ 

5. Let *Y* be a random variable with pdf given by  $f(y) = \begin{cases} \frac{y}{2} & \text{if } 0 \le y \le 2, \\ 0 & \text{otherwise.} \end{cases}$ 

(a) Find a value  $\eta_{0.25}$  such that  $P(Y \le \eta_{0.25}) = 0.25$ .



(b) What is the median of *Y*?

