

EXAM 1:

A: 63 - 70
 B: 54 - 62
 C: 40 - 53
 D: below 40

Questions about the exam?
 Talk with Prof. Wright!

From last time:

3. An unknown number, N , of animals inhabit a certain region. To estimate the size of the population, ecologists perform the following experiment: They first catch M of these animals, mark them in some way, and release them. After allowing the animals to disperse throughout the region, they catch n of the animals and count the number, X , of marked animals in this second catch.

The ecologists want to make a *maximum likelihood estimate* of the population size N . This means that if the observed value of X is x , then they estimate the population size to be the integer N that maximizes the probability that $X = x$. Help them complete this estimate as follows.

$X \sim \text{Hypergeometric}(n, M, N)$

Let $P_x(N)$ be the probability that $X = x$ given that N is the true population size.

So:
$$P_x(N) = \frac{\binom{M}{x} \binom{N-M}{n-x}}{\binom{N}{n}} \quad \text{and} \quad \frac{P_x(N)}{P_x(N-1)} = \frac{(N-M)(N-n)}{N(N+x-M-n)}$$

$$\frac{P_x(N)}{P_x(N-1)} \geq 1 \quad \text{iff} \quad \frac{(N-M)(N-n)}{N(N+x-M-n)} \geq 1$$

$$\text{iff} \quad (N-M)(N-n) \geq N(N+x-M-n)$$


$$\text{iff} \quad \cancel{N^2 - Mn} - \cancel{Nn} + Mn \geq \cancel{N^2} + Nx - \cancel{NM} - \cancel{Nn}$$

$$\text{iff} \quad Mn \geq Nx$$

$$\text{iff} \quad \frac{Mn}{x} \geq N$$

If $N \leq \frac{Mn}{x}$, then $P_x(N) \geq P_x(N-1)$, so population size N is more likely than population size $N-1$.

If $N > \frac{Mn}{x}$ then population size N is less likely than $N-1$.



The most likely population is the largest integer N that is less than or equal to $\frac{Mn}{x}$.

(f) If $M=30$, $n=20$, $x=7$,
 then max. likelihood estimate for N is: $\frac{Mn}{x} = \frac{30(20)}{7} \approx 85.7$
 so $N=85$.

NEGATIVE BINOMIAL DISTRIBUTION

An experiment consists of a sequence of independent trials. Each trial results in either "success" or "failure." The probability of success is p for each trial. The experiment stops when a certain number, r , of successes have occurred. Let X be the number of trials necessary to achieve r successes.

Then $X \sim \text{Negative Binomial}(r, p)$

$$\text{pmf: } P(X=x) = \binom{x-1}{r-1} p^r (1-p)^{x-r}$$

If $r=1$, then $X \sim \text{Geometric}(p)$

$$\text{pmf: } P(X=x) = (1-p)^{x-1} p$$